

p16 Monoclonal antibody

Cat:B32160R

Company: Haokebio

Uniprot ID:P42771

Applications: IHC:1:100-1:200

Organism:Mouse

IHC-Polymer:1:400-1:800

Species reactivity:Human

TSA:1:500-1:1000

Background:


Steroidogenic Factor 1 (SF-1)

4°C overnight.Secondary antibody: Poly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

Protein full name:

cyclin-dependent kinase inhibitor 2A

Images:



Synonyms:

P16, p16 INK4, P16 INK4A, p16INK4A, P16-INK4A

Immunogen:

Peptide

Isotype:

IgG

Subcellular location:

Nucleus

Sample: Human Cervical intraepithelial neoplasia tissue,4% PFA 12-24h

Purity:

Affinity purification

Source of Reagents:

发表[中文论文]请标注:p16(B32160R)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:p16(B32160R) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.

Form:

Liquid

Storage Buffer:

PBS with 0.02%sodium azide,100 µg/ml BSA and 50% glycerol.

Storage:

Store at -20 °C for one year.

Experimental procedure:

Antigen retrieval: Citrate buffer (pH 9.0) , Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes.Incubate antibody,