

## MHC Class I Recombinant antibody

**Cat:** B35002D

**Company:** HaoKebio

**Uniprot ID:** P01889

**Applications:** IHC:1:500-1:2000

**Organism:** Rabbit

IHC-Polymer:1:2000-1:8000

**Species reactivity:** Human

IHC-TSA:1:2500-1:10000

**Molecular Weight Calculation:** 41 kDa

IF:1:50-1:100

**Observed Molecular Weight:** 41 kDa

WB:1:1000

### Background:

HLA-A belongs to the HLA class I heavy chain paralogues. This class I molecule is a heterodimer consisting of a heavy chain and a light chain (beta-2 microglobulin). The heavy chain is anchored in the membrane. Class I molecules play a central role in the immune system by presenting peptides derived from the endoplasmic reticulum lumen. They are expressed in nearly all cells. The heavy chain is approximately 45 kDa and its gene contains 8 exons. Exon 1 encodes the leader peptide, exons 2 and 3 encode the alpha1 and alpha2 domains, which both bind the peptide, exon 4 encodes the alpha3 domain, exon 5 encodes the transmembrane region, and exons 6 and 7 encode the cytoplasmic tail. Polymorphisms within exon 2 and exon 3 are responsible for the peptide binding specificity of each class one molecule. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. Hundreds of HLA-A alleles have been described.

### Protein full name:

Major Histocompatibility Complex Class I Molecule

### Synonyms:

HLAA

### Immunogen:

Recombinant protein

### Isotype:

IgG

### Subcellular location:

Membrane

### Purity:

Affinity purification

### Form:

Liquid

### Storage Buffer:

PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.


### Storage:

Store at -20 °C for one year.

### Experimental procedure:

Antigen retrieval: Citrate buffer (pH 9.0), Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: Poly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

### Images:



Dilution of 1:1000 incubated at room temperature for 1.5 hours.

### Source of Reagents:

发表[中文论文]请标注:MHC Class I(B35052D)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:MHC Class I(B35052D) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.

